



CERTIFICATION

AOAC Research Institute
Performance Tested MethodsSM

Certificate No.
081904

The AOAC Research Institute hereby certifies the method known as:

iQ-Check S. Typhimurium Real-Time PCR

Corporate Location
Bio-Rad Laboratories
2000 Alfred Nobel Drive
Hercules, CA 94547 USA

Manufacturing Location Bio-Rad
Laboratories
925 Alfred Nobel Drive
Hercules, CA 94547 USA

This method has been evaluated in the AOAC Research Institute *Performance Tested MethodsSM* Program and found to perform as stated in the applicability of the method. This certificate indicates an AOAC Research Institute Certification Mark License Agreement has been executed which authorizes the manufacturer to display the AOAC Research Institute *Performance Tested MethodsSM* certification mark on the above-mentioned method for the period below. Renewal may be granted by the Expiration Date under the rules stated in the licensing agreement.

Scott Coates

Scott Coates, Senior Director
Signature for AOAC Research Institute

Issue Date	October 30, 2023
Expiration Date	December 31, 2024

AUTHORS ORIGINAL VALIDATION: Mike Clark, Jean-Philippe Tourniaire, Sophie Pierre, Christophe Quiring MODIFICATION JANUARY 2023: Mike Clark	SUBMITTING COMPANY Bio-Rad Laboratories 2000 Alfred Nobel Drive Hercules, CA 94547 USA
METHOD NAME iQ-Check S. Typhimurium Real-Time PCR	CATALOG NUMBER 12004306
INDEPENDENT LABORATORY Q Laboratories 1930 Radcliffe Dr. Cincinnati, OH USA 45214	
APPLICABILITY OF METHOD Target Analyte – <i>Salmonella</i> Typhimurium.	REFERENCE METHODS U.S. Department of Agriculture Food Safety and Inspection Service (2018) Microbiology Laboratory Guidebook, Revision 4.09 <i>Isolation and Identification of Salmonella from Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges</i> (2) FDA, <i>Environmental Sampling and Detection of Salmonella in Poultry Houses</i> . (3)
Matrixes – (25 g) – raw chicken breast with skin, raw chicken breast without skin, raw chicken breast containing 2% w/w salt, raw chicken thigh with skin, raw chicken thigh without skin Environmental boot swabs, drag swabs	
Performance claims – The study data detected no statistical difference between the iQ-Check S. Typhimurium method and the reference methods.	
ORIGINAL CERTIFICATION DATE August 21, 2019	CERTIFICATION RENEWAL RECORD Renewed annually through December 2024.
METHOD MODIFICATION RECORD 1. January 2020 Level 1 2. January 2021 Level 1 3. April 2021 Level 1 4. October 2021 Level 1 5. January 2023 Level 2 6. October 2023 Level 1	SUMMARY OF MODIFICATION 1. Insert reformatted. 2. Editorial/clerical changes. 3. Software was updated from Version 3 to Version 4 allowing compatibility with Windows 10. 4. Editorial changes and addition of user information in French, German, Spanish, Portuguese, and Italian. 5. Addition of CFX Opus Deepwell, with CFX Manager Software, Industrial Diagnostic Edition version 3.1 using Free DNA Removal Solution and Fast APF protocols. 6. Editorial/clerical changes.
Under this AOAC Performance Tested Methods SM License Number, 081904 this method is distributed by: NONE	Under this AOAC Performance Tested Methods SM License Number, 081904 this method is distributed as: NONE
PRINCIPLE OF THE METHOD (1) The iQ-Check S. Typhimurium real-time PCR assay has the flexibility to be used in multiple ways. The kit can be used as a primary screening tool for the direct detection of <i>S. Typhimurium</i> (ST) in select foods and primary production samples such as boot or drag swabs. The kit can also be used as a secondary screening tool after first screening for presumptive positive samples with the iQ-Check <i>Salmonella</i> spp. kit using the same extracted DNA enabling quicker confirmation of ST present in a sample. Finally, the kits can be used to identify isolated suspected ST colonies directly from selective agar plates such as RAPID'Salmonella chromogenic agar. The iQ-Check S. Typhimurium real-time PCR assay is based on gene amplification and detection by real-time polymerase chain reaction (PCR) technology. Ready-to-use PCR reagents contain oligonucleotides (primers and probes) specific for <i>S. Typhimurium</i> , as well as DNA polymerase and nucleotides. Detection and data analysis are optimized for use with Bio-Rad real-time PCR instruments, such as the CFX 96 Touch Deep Well system. The iQ-Check Prep is a robotic liquid handling platform that performs DNA extraction and PCR plate set-up. It is designed for all the iQ-Check Food Pathogen Detection Kits and the CFX96 Touch Deep Well real-time PCR System. It is a completely integrated automated solution for food pathogen testing. The iQ-Check Free DNA Removal Solution is also provided in a separate kit to safely reduce free DNA present in the matrixes. PCR is a powerful technique used to generate many copies of target DNA. During the PCR reaction, several cycles of heating and cooling facilitate DNA denaturation, primer binding to the target region, and DNA polymerase extension of the DNA, creating copies of the target region. These copies are called amplicons. A synthetic DNA "internal control" is included in the reaction mix. This control is amplified with a specific probe at the same time as the target analytes. It allows for the validation of any negative result (4).	

DISCUSSION OF THE VALIDATION STUDY (1)

The iQ-Check S. Typhimurium Assays successfully detected ST from all six matrixes analyzed. The ability to screen for ST allows the end user to follow the current guidelines outlined by the USDA/FSIS MLG 4.09 and FDA Env Sampling & Det of *Salm* in Poultry Houses and identify two of the most prevalent serotypes in the poultry industry with two assays. The alternative confirmation procedure allows the end user to confirm samples at the same degree of sensitivity as the USDA/FSIS MLG 4.09 and FDA Env Sampling & Det of *Salm* in Poultry Houses, but one to two days sooner than the traditional confirmation method. With the addition of the Free DNA Removal Solution, the test kits allow the user to safely reduce free DNA present in the matrixes.

In the inclusivity and exclusivity evaluations, all inclusivity organisms were correctly identified, and all exclusivity organisms were correctly excluded. The inclusivity data also demonstrated that positive isolated colonies from the RAPID' *Salmonella* could be tested on the kit as part of the overall confirmation protocol for ST. In the method comparison study, the iQ-Check test kit demonstrated no statistically significant difference between candidate and reference method results (dPOD_C), or between presumptive and confirmed results (dPOD_{CF}) for all target pathogens.

The lot-to-lot consistency and stability study show no significant differences observed across the shelf life of the kits for three different lots of kits at each time point tested. The robustness study shows no discrepant results observed for all 8 treatment combinations for the *S. Typhimurium* assay.

The iQ-Check S. Typhimurium Assay is quick and simple to perform, providing results in approximately 4 hours post incubation of the enrichment for up to 94 sample replicates. The use of the iQ-Check Prep Instrument provides a hands-free application that can reduce possible contamination caused by the analyst performing testing. The iQ-Check Prep Instrument can perform DNA extraction and PCR preparation and provides added value of traceability to the lab. The CFX Manager IDE software is user friendly with the ability to track lot information and sample identification quickly and with ease. Because results are displayed in real-time, the user is able to quickly and accurately determine if results will be valid before the end of the run. The software also provides the user the option to analyze each individual Cq curves to help aid in problem solving any issues within an individual reaction.

Table 3. Inclusivity Results for iQ-Check S. Typhimurium Assay (1)

No.	Species	Subspecies	Serovar	Antigenic Formula	Source	Origin	BPW	BPW + Suppl.	TCS	RSA
1	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	¹ ATCC 14028	Chicken heart and liver tissue	+	+	+	+
2	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	² n° Anses: 38.09	Brine	+	+	+	+
3	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 442.09	Beef meat	+	+	+	+
4	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 447.09	Pork (crépine)	+	+	+	+
5	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 591.09	Lamb with sauce	+	+	+	+
6	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 695.09	Stuffed quail	+	+	+	+
7	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 839.09	Culture from veal feces	+	+	+	+
8	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 838.09	Environment (Duck)	+	+	+	+
9	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 553.11	Culture from horse feces	+	+	+	+
10	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 564.11	White pepper	+	+	+	+
11	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 708.11	Hoki fillet with cream	+	+	+	+
12	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 781.11	Pigeon viscera	+	+	+	+
13	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 792.11	Urban compost	+	+	+	+
14	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 835.11	Env. (Chicken)	+	+	+	+
15	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 840.11	Streaky ham	+	+	+	+
16	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 845.11	Whole quail	+	+	+	+
17	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 880.11	Spareribs	+	+	+	+
18	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 886.11	Culture from swine feces	+	+	+	+
19	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 907.11	Raw milk (cow)	+	+	+	+
20	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 976.11	Sausage	+	+	+	+
21	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 977.11	Tomato filling	+	+	+	+
22	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 979.11	Stuffed potatoes	+	+	+	+
23	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 985.11	Fish meal	+	+	+	+
24	<i>Salmonella</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses:	Pet food	+	+	+	+

	<i>enterica</i>				1175.11				
25	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	n° Anses: 119.11	Foie gras (liver)	+	+	+
26	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	³ ADRIA n°4	Milk powder	+	+	+
27	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°13	Pasteurized liquid egg	+	+	+
28	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°206	Pasteurized liquid egg	+	+	+
29	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°472	Egg yolk	+	+	+
30	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°776	Pasteurized liquid egg	+	+	+
31	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	⁴ CIP 58.58	Ready-to-eat	+	+	+
32	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°19	Foie (Liver)	+	+	+
33	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°22	Raw ground meat	+	+	+
34	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°167	Ready-to-eat	+	+	+
35	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°193	Chipolatas sausages	+	+	+
36	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°830	Chipolatas sausages	+	+	+
37	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°911	Merguez sausages	+	+	+
38	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°987	Chipolatas sausages	+	+	+
39	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°4874	Meat (pâté)	+	+	+
40	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°A00C003	Frozen meat	+	+	+
41	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°A00C059	Frozen beef trim	+	+	+
42	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°A00C060	ground beef	+	+	+
43	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n°Ad1070	Pork	+	+	+
44	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° ST325	Pork	+	+	+
45	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° ST1	Pork	+	+	+
46	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° ST394	Pork	+	+	+
47	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° ST719	Pork	+	+	+
48	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° ST11	Pork	+	+	+
49	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° JES411	Liquid egg	+	+	+
50	<i>Salmonella enterica</i>	<i>enterica</i>	Typhimurium	1,4,[5],12:i:1,2	ADRIA n° Ad913	Beef trim	+	+	+

¹American Type Culture Collection, Manassas, VA²French Agency for Food, Environmental and Occupational Health & Safety, Maisons-Alfort, France³Culture Collection ADRIA Developpement, Quimper, France⁴Collection Institute Pasteur, Paris, France

Table 4. Exclusivity Results for iQ-Check S. Typhimurium Assay (1)

No.	Species	Subspecies	Serovar	Antigenic Formula	Source	Origin	BPW	BPW + Suppl.	TCS	RSA
1	<i>Salmonella bongori</i>			V 66:z41:-	¹ CIP 82.33T	Unknown	-	-	-	-
2	<i>Salmonella bongori</i>			V 48:z35:-	² RDCM 001	Unknown	-	-	-	-
3	<i>Salmonella enterica</i>	arizonaee		51 : z4,z23	CIP 82.30T	Unknown	-	-	-	-
4	<i>Salmonella enterica</i>	arizonaee			³ ADRIA/AFSSA n° Ad 478	Shell fish	-	-	-	-
5	<i>Salmonella enterica</i>	arizonaee			⁴ RDC 36	Unknown	-	-	-	-
6	<i>Salmonella enterica</i>	arizonaee			ADRIA/AFSSA n° Ad 450	Sheep milk	-	-	-	-
7	<i>Salmonella enterica</i>	arizonaee			ADRIA/AFSSA n° Ad 451	Unknown	-	-	-	-
8	<i>Salmonella enterica</i>	arizonaee			ADRIA/AFSSA n° Ad 452	Unknown	-	-	-	-
9	<i>Salmonella enterica</i>	arizonaee			ADRIA/AFSSA n° Ad 453	Unknown	-	-	-	-
10	<i>Salmonella enterica</i>	arizonaee			RDM 1 Hollande	Unknown	-	-	-	-
11	<i>Salmonella enterica</i>	diarizonae		IIIb 6,7 : l,v:z53	CIP 82.31T	Unknown	-	-	-	-
12	<i>Salmonella enterica</i>	diarizonae		IIIb 38 : l,v:z35	ADRIA/AFSSA n°Ad 594	Frog legs	-	-	-	-
13	<i>Salmonella enterica</i>	diarizonae		IIIb 61 : i:z53	ADRIA/AFSSA n° Ad 595	Cheese	-	-	-	-
14	<i>Salmonella enterica</i>	enterica	Abaetetuba	11 : k:1,5	⁵ ATCC 35640	Unknown	-	-	-	-
15	<i>Salmonella enterica</i>	enterica	Aberdeen	11 : i:1,2	⁶ CMF 114.1	Unknown	-	-	-	-
16	<i>Salmonella enterica</i>	enterica	Abony	1,4,12,27:b:e,n,x	⁷ NCTC 6017	Unknown	-	-	-	-
17	<i>Salmonella enterica</i>	enterica	Adelaïde	35 : f,g : -	CMF 482.2	Unknown	-	-	-	-
18	<i>Salmonella enterica</i>	enterica	Agama	4,12 : i:1,6	RDCM 205 Hey's food (NL)	Unknown	-	-	-	-
19	<i>Salmonella enterica</i>	enterica	Agona	1,4,[5],12 : f,g,s:[1,2]	RDCM 057	Unknown	-	-	-	-
20	<i>Salmonella enterica</i>	enterica	Albany	8,20 : z4,z24:-	CMF 82.2	Unknown	-	-	-	-
21	<i>Salmonella enterica</i>	enterica	Anatum	3,{10}{15}{15,34} : e,h:1,6	CMF 814.4	Unknown	-	-	-	-
22	<i>Salmonella enterica</i>	enterica	Bambylor	9,46 : z:e,n,z15	CMF 135.1	Unknown	-	-	-	-
23	<i>Salmonella enterica</i>	enterica	Bareilly	6,7,14 : y:1,5	CMF 136.1	Unknown	-	-	-	-
24	<i>Salmonella enterica</i>	enterica	Berta	1,9,12 : [f],g,[t]:-	CMF 141.2	Unknown	-	-	-	-
25	<i>Salmonella enterica</i>	enterica	Blegdam	9,12 : g,m,q:-	NCTC 5769	Unknown	-	-	-	-
26	<i>Salmonella enterica</i>	enterica	Blockley	6,8 : k:1,5	RDCM 127 ADRIA	Unknown	-	-	-	-
27	<i>Salmonella enterica</i>	enterica	Braenderup	6,7,14 : e,h:e,n,z15	CMF 151	Unknown	-	-	-	-
28	<i>Salmonella enterica</i>	enterica	Brandenburg	4,[5],12 : l,v:e,n,z15	RDC 106	Unknown	-	-	-	-
29	<i>Salmonella enterica</i>	enterica	Bredeney	1,4,12,27 : l,v:1,7	CMF 817.1	Unknown	-	-	-	-
30	<i>Salmonella enterica</i>	enterica	California	4,12 : g,m,t:[z67]	RDCM 051	Unknown	-	-	-	-
31	<i>Salmonella enterica</i>	enterica	Cerro	6,14,18 : z4,z23:[1,5]	CMF 166.2	Unknown	-	-	-	-
32	<i>Salmonella enterica</i>	enterica	Carrau	6,14,[24] : y:1,7	CMF 142.2	Unknown	-	-	-	-
33	<i>Salmonella enterica</i>	enterica	Crossness	67 : r:1,2	CMF 165.1	Unknown	-	-	-	-
34	<i>Salmonella enterica</i>	enterica	Cubana	1,13,23 : z29:-	CMF 188	Unknown	-	-	-	-

35	<i>Salmonella enterica</i>	<i>enterica</i>	Choleraesuis	6,7 : c:1,5	CMF 181.1	Unknown	-	-	-	-
36	<i>Salmonella enterica</i>	<i>enterica</i>	Dahlem	48 : k:e,n,z15	CMF 924.1	Unknown	-	-	-	-
37	<i>Salmonella enterica</i>	<i>enterica</i>	Derby	1,4,[5],12 : f,g:[1,2]	RDC 91	Unknown	-	-	-	-
38	<i>Salmonella enterica</i>	<i>enterica</i>	Dublin	1,9,12[Vi] : g,p:-	RDC 31	Unknown	-	-	-	-
39	<i>Salmonella enterica</i>	<i>enterica</i>	Duisburg	1,4,12,[27] : d:e,n,z15	RDCM 199 Hey's food (NL)	Unknown	-	-	-	-
40	<i>Salmonella enterica</i>	<i>enterica</i>	Emek	8,20 : g,m,s:-	ADRIA n° Ad333	Unknown	-	-	-	-
41	<i>Salmonella enterica</i>	<i>enterica</i>	Enteritidis	1,9,12 : g,m:-	RDC 338	Unknown	-	-	-	-
42	<i>Salmonella enterica</i>	<i>enterica</i>	Fischerkietz	1,6,14,25 : y:e,n,x	RDCM 202 Hey's food (NL)	Unknown	-	-	-	-
43	<i>Salmonella enterica</i>	<i>enterica</i>	Ferruch	8 : e,h:1,5	RDCM 207 Hey's food (NL)	Unknown	-	-	-	-
44	<i>Salmonella enterica</i>	<i>enterica</i>	Gallinarum	9,12:-:-	⁸ DSM 13674	Unknown	-	-	-	-
45	<i>Salmonella enterica</i>	<i>enterica</i>	Gaminara	16 : d:1,7	CMF 221.1	Unknown	-	-	-	-
46	<i>Salmonella enterica</i>	<i>enterica</i>	Give	3,{10}{15}{15,34} : l,v:1,7	RDM 2 Hollande	Unknown	-	-	-	-
47	<i>Salmonella enterica</i>	<i>enterica</i>	Glostrup	6,8 : z10:e,n,z15	CMF 226.3	Unknown	-	-	-	-
48	<i>Salmonella enterica</i>	<i>enterica</i>	Goldcoast	6,8 : r:l,w	RDCM 210 Hey's food (NL)	Unknown	-	-	-	-
49	<i>Salmonella enterica</i>	<i>enterica</i>	Grumpensis	1,13,23 : d:1,7	CMF 478.2	Unknown	-	-	-	-
50	<i>Salmonella enterica</i>	<i>enterica</i>	Guinea	1,44 : z10:1,7	RDCM 062	Unknown	-	-	-	-
51	<i>Salmonella enterica</i>	<i>enterica</i>	Hadar	6,8 : z10:e,n,x	CMF 234.1	Unknown	-	-	-	-
52	<i>Salmonella enterica</i>	<i>enterica</i>	Havana	1,13,23 : f,g,[s]:-	CMF 237.1	Unknown	-	-	-	-
53	<i>Salmonella enterica</i>	<i>enterica</i>	Heidelberg	1,4,[5],12 : r:1,2	RDC 93	Unknown	-	-	-	-
54	<i>Salmonella enterica</i>	<i>enterica</i>	Illinois	3,{10}{15}{15,34} : z10:1,5	CMF 251.2	Unknown	-	-	-	-
55	<i>Salmonella enterica</i>	<i>enterica</i>	Indiana	1,4,12 : z:1,7	RDCM 128 ADRIA	Unknown	-	-	-	-
56	<i>Salmonella enterica</i>	<i>enterica</i>	Infantis	6,7,14 : r:1,5	CIP 82.97	Unknown	-	-	-	-
57	<i>Salmonella enterica</i>	<i>enterica</i>	Inverness	38 : k:1,6	CMF 253.1	Unknown	-	-	-	-
58	<i>Salmonella enterica</i>	<i>enterica</i>	Johannesburg	1,40 : b:e,n,x	CMF 256.1	Unknown	-	-	-	-
59	<i>Salmonella enterica</i>	<i>enterica</i>	Kedougou	1,13,23 : i:l,w	RDCM 212 Hey's food (NL)	Unknown	-	-	-	-
60	<i>Salmonella enterica</i>	<i>enterica</i>	Kentucky	8,20 : i:z6	CMF 264.2	Unknown	-	-	-	-
61	<i>Salmonella enterica</i>	<i>enterica</i>	Kottbus	6,8 : e,h:1,5	RDCM 198 Hey's food (NL)	Unknown	-	-	-	-
62	<i>Salmonella enterica</i>	<i>enterica</i>	Livingstone	6,7,14 : d:l,w	RDCM 155 Friesland Campina	16S ID	-	-	-	-
63	<i>Salmonella enterica</i>	<i>enterica</i>	Livingstone	6,7,14 : d:l,w	⁹ LAV n°1	Env.	-	-	-	-
64	<i>Salmonella enterica</i>	<i>enterica</i>	Lomita	6,7 : e,h:1,5	CMF 125.3	Unknown	-	-	-	-
65	<i>Salmonella enterica</i>	<i>enterica</i>	London	3,{10}{15} : l,v:1,6	RDCM 129 ADRIA	Unknown	-	-	-	-
66	<i>Salmonella enterica</i>	<i>enterica</i>	Mbandaka	6,7,14 : z10:e,n,z15	RDCM 130 ADRIA	Unknown	-	-	-	-
67	<i>Salmonella enterica</i>	<i>enterica</i>	Miami	1,9,12 : a:1,5	CMF 307	Unknown	-	-	-	-
68	<i>Salmonella enterica</i>	<i>enterica</i>	Minnesota	21 : b:e,n,x	CMF 146.3	Unknown	-	-	-	-
69	<i>Salmonella enterica</i>	<i>enterica</i>	Montevideo	6,7,14,[54] : g,m,[p],s:[1,2,7]	RDCM 45	Unknown	-	-	-	-

70	<i>Salmonella enterica</i>	<i>enterica</i>	Moscow	1,9,12 : g,q:-	RDCM 053	Unknown	-	-	-	-
71	<i>Salmonella enterica</i>	<i>enterica</i>	Muenchen	6,8 : d:1,2	CMF 337	Unknown	-	-	-	-
72	<i>Salmonella enterica</i>	<i>enterica</i>	Newport	6,8,20 : e,h:1,2	RDCM 208 Hey's food (NL)	Unknown	-	-	-	-
73	<i>Salmonella enterica</i>	<i>enterica</i>	Nienstedten	6,7,14 : b:l,w	CMF 352	Unknown	-	-	-	-
74	<i>Salmonella enterica</i>	<i>enterica</i>	Nottingham	16 : d:e,n,z15	NCTC 7832T	Unknown	-	-	-	-
75	<i>Salmonella enterica</i>	<i>enterica</i>	Ohio	6,7,14 : b:l,w	RDCM 132 ADRIA	Unknown	-	-	-	-
76	<i>Salmonella enterica</i>	<i>enterica</i>	Oranienburg	6,7,14 : m,t: [z57]	CMF 360	Unknown	-	-	-	-
77	<i>Salmonella enterica</i>	<i>enterica</i>	Ouakam	9,46 : z29:-	CMF 364	Unknown	-	-	-	-
78	<i>Salmonella enterica</i>	<i>enterica</i>	Paratyphi B Java	1,4,[5],12 : b:1,2	RDCM 192 Hey's food (NL)	Unknown	-	-	-	-
79	<i>Salmonella enterica</i>	<i>enterica</i>	Potsdam	6,7,14 : l,v:e,n,z15	CMF 225.2	Unknown	-	-	-	-
80	<i>Salmonella enterica</i>	<i>enterica</i>	Putten	13,23 : d:l,w	RDCM 214	Unknown	-	-	-	-
81	<i>Salmonella enterica</i>	<i>enterica</i>	Rostock	1,9,12:g,p,u:-	RDCM 55	Unknown	-	-	-	-
82	<i>Salmonella enterica</i>	<i>enterica</i>	Rubislaw	11 : r:e,n,x	CMF 414.2	Unknown	-	-	-	-
83	<i>Salmonella enterica</i>	<i>enterica</i>	Saintpaul	1,4,[5],12 : e,h:1,2	RDCM 101 ADRIA	Ad203	-	-	-	-
84	<i>Salmonella enterica</i>	<i>enterica</i>	Senftenberg	1,3,19 : g,[s],t:-	RDCM 209 Hey's food (NL)	Unknown	-	-	-	-
85	<i>Salmonella enterica</i>	<i>enterica</i>	Sundsvall	[1],6,14,[25] : z:e,n,x	CMF 877.2	Unknown	-	-	-	-
86	<i>Salmonella enterica</i>	<i>enterica</i>	Taksony	1,3,19 : i:z6	RDCM 203 Hey's food (NL)	Unknown	-	-	-	-
87	<i>Salmonella enterica</i>	<i>enterica</i>	Tallahassee	6,8 : z4,z32:-	CMF 822.1	Unknown	-	-	-	-
88	<i>Salmonella enterica</i>	<i>enterica</i>	Tennessee	6,7,14 : z29:[1,2,7]	RDCM Nestlé 599	Env.	-	-	-	-
89	<i>Salmonella enterica</i>	<i>enterica</i>	Thompson	6,7,14 : k:1,5	RDCM 056	Unknown	-	-	-	-
90	<i>Salmonella enterica</i>	<i>enterica</i>	Tournai	3,{10}{15} : y:z6	CMF 448.1	Unknown	-	-	-	-
91	<i>Salmonella enterica</i>	<i>enterica</i>	Virchow	6,7,14 : r:1,2	CMF 805.2	Unknown	-	-	-	-
92	<i>Salmonella enterica</i>	<i>enterica</i>	Wayne	30:g,z51:-	RDCM 54	Unknown	-	-	-	-
93	<i>Salmonella enterica</i>	<i>enterica</i>	Worthington	1,13,23 : z :l,w	RDCM 157 Friesland Campina	16S ID	-	-	-	-
94	<i>Salmonella enterica</i>	<i>enterica</i>	Yoruba	16 : c:l,w	CMF 3913/83	Unknown	-	-	-	-
95	<i>Salmonella enterica</i>	<i>houtenae</i>		IV 45 : g,z51:-	CIP 82.32T	Unknown	-	-	-	-
96	<i>Salmonella enterica</i>	<i>houtenae</i>		IV 50 : g,z51:-	ADRIA/AFSSA n° Ad 596	Milk product	-	-	-	-
97	<i>Salmonella enterica</i>	<i>houtenae</i>		IV 43 : z4,z32:-	ADRIA/AFSSA n° Ad 597	Cooked fish	-	-	-	-
98	<i>Salmonella enterica</i>	<i>indica</i>		VI [1],6,14,[25] : a:e,n,x	CIP 102501T	Unknown	-	-	-	-
99	<i>Salmonella enterica</i>	<i>salamae</i>	Betioky	II 59 : k:z65	CMF 141.1	Unknown	-	-	-	-
100	<i>Salmonella enterica</i>	<i>salamae</i>	Grabouw	II 11 : g,m,s,t:z39	RDCM 047	Unknown	-	-	-	-
101	<i>Salmonella enterica</i>	<i>salamae</i>	Manica	II 1,9,12 : g,m,s,t:(z42)	RDCM 048	Unknown	-	-	-	-
102	<i>Salmonella enterica</i>	<i>salamae</i>	Daressalaam	II 1,9,12 : l,w:e,n,x	CIP 82.29T	Unknown	-	-	-	-
103	<i>Salmonella enterica</i>	<i>salamae</i>		II 42 : g,t:-	ADRIA/AFSSA n° Ad 592	Kangaroo meat	-	-	-	-
104	<i>Salmonella enterica</i>	<i>salamae</i>		II 42 : b:e,n,x,z15	ADRIA/AFSSA n° Ad 593	Seed	-	-	-	-

105	<i>Salmonella enterica</i>	<i>salamae</i>	Tranoroa	II 55 : k:z39	CMF 463.1	Unknown	-	-	-	-
106	<i>Salmonella enterica</i>	<i>salamae</i>	Zuerich	II 1,9,12,46,27 : c:z39	CMF 510.1	Unknown	-	-	-	-
107	<i>Acinetobacter baumanii</i>				RDC 146	Unknown	-		-	
108	<i>Aeromonas hydrophila</i>				¹⁰ LMG 2844T	Frog red-leg	-		-	
109	<i>Aeromonas hydrophila/caviae</i>				RDCM 152 Friesland Campina	16S ID	-		-	
110	<i>Bacillus licheniformis</i>				RDC 88	Unknown	-		-	
111	<i>Bacillus cereus</i>				RDCM 218 Friesland Campina	16S ID	-		-	
112	<i>Campylobacter jejuni</i>				ATCC 29428	Unknown	-		-	
113	<i>Campylobacter coli</i>				ATCC 43489	Unknown	-		-	
114	<i>Campylobacter lari</i>				ATCC 35221	Unknown	-		-	
115	<i>Campylobacter upsaliensis</i>				RDC 68	Unknown	-		-	
116	<i>Citrobacter freundii</i>				RDCM 154 Friesland Campina	16S ID	-		-	
117	<i>Citrobacter freundii</i>				ATCC 8090	Unknown	-		-	
118	<i>Cronobacter sakazakii</i>				RDCM 210 Friesland Campina	16S ID	-		-	
119	<i>Cronobacter sakazakii</i>				RDCM 213 Friesland Campina	16S ID	-		-	
120	<i>Enterobacter aerogenes</i>				ATCC 13048	Unknown	-		-	
121	<i>Enterobacter amnigenus</i>				RDCM 204 Friesland Campina	16S ID	-		-	
122	<i>Enterobacter amnigenus</i>				RDCM 144 Friesland Campina	16S ID	-		-	
123	<i>Enterobacter asburiae</i>				RDCM 142 Friesland Campina	16S ID	-		-	
124	<i>Enterobacter cloaceae</i>				RDCM 151 Friesland Campina	16S ID	-		-	
125	<i>Enterobacter cloaceae</i>				LMG 2783	Unknown	-		-	
126	<i>Enterobacter cowanii</i>				RDCM 202 Friesland Campina	16S ID	-		-	
127	<i>Enterobacter pyrinus</i>				RDCM 211 Friesland Campina	16S ID	-		-	
128	<i>Enterobacter sakazakii</i>				RDC 236	Milk product	-		-	
129	<i>Enterobacter sakazakii</i>				RDCM 150 Friesland Campina	16S ID	-		-	
130	<i>Enterococcus faecium</i>				Bio-Rad RIVH WR63	Unknown	-		-	
131	<i>Escherichia coli</i>			O157H7 VT-	ATCC 700728	Unknown	-		-	
132	<i>Escherichia coli</i>				RDCM 203 Friesland Campina	16S ID	-		-	
133	<i>Escherichia hermanii</i>				RDC 72 ADRIA ADRIA	White egg	-		-	
134	<i>Hafnia alvei</i>				CIP 57.31T	Unknown	-		-	

135	<i>Hafnia alvei</i>				RDCM 145 Friesland Campina	16S ID	-		-	
136	<i>Hafnia alvei</i>				RDCM 205 Friesland Campina	16S ID	-		-	
137	<i>Klebsiella oxytoca</i>				RDC 30	Unknown	-		-	
138	<i>Klebsiella oxytoca</i>				RDCM 208 Friesland Campina	16S ID	-		-	
139	<i>Klebsiella oxytoca</i>				RDCM 148 Friesland Campina	16S ID	-		-	
140	<i>Klebsiella pneumoniae</i>				RDCM 209 Friesland Campina	16S ID	-		-	
141	<i>Klebsiella pneumoniae</i>	<i>pneumoniae</i>			RDCM 149 Friesland Campina	16S ID	-		-	
142	<i>Listeria monocytogenes</i>			1/2b	¹¹ CNRL 87373	Unknown	-		-	
143	<i>Listeria monocytogenes</i>			1/2c	CNRL 103573	Unknown	-		-	
144	<i>Listeria monocytogenes</i>			1/2a	CIP 78.31	mesenteric lymph node, guinea pig	-		-	
145	<i>Micrococcus luteus</i>				RDC 70	soil	-		-	
146	<i>Pantoea agglomerans</i>				RDCM 141 Friesland Campina	16S ID	-		-	
147	<i>Pantoea agglomerans</i>				RDCM 201 Friesland Campina	16S ID	-		-	
148	<i>Proteus mirabilis</i>				ATCC 29906	Unknown	-		-	
149	<i>Proteus mirabilis</i>				ATCC 25933	Unknown	-		-	
150	<i>Proteus mirabilis</i>				RDCM T04.37.17	Roquefort	-		-	
151	<i>Pseudomonas aeruginosa</i>				RDCM 153 Friesland Campina	16S ID	-		-	
152	<i>Pseudomonas aeruginosa</i>				RDCM 212 Friesland Campina	16S ID	-		-	
153	<i>Raoultella terrigena</i>				RDCM 146 Friesland Campina	16S ID	-		-	
154	<i>Raoultella terrigena</i>				RDCM 206 Friesland Campina	16S ID	-		-	
155	<i>Serratia marcescens</i>				ATCC 8100	Unknown	-		-	
156	<i>Serratia marcescens</i>				RDCM 147 Friesland Campina	16S ID	-		-	
157	<i>Serratia marcescens</i>				RDCM 207 Friesland Campina	16S ID	-		-	
158	<i>Shigella flexneri</i>				ATCC 12022	Unknown	-		-	
159	<i>Shigella sonnei</i>				ATCC 25931	Human feces	-		-	
160	<i>Staphylococcus aureus</i>				ATCC 25923	Unknown	-		-	
161	<i>Staphylococcus aureus</i>				RDCM 217 Friesland Campina	16S ID	-		-	
162	<i>Staphylococcus epidermidis</i>				RDCM 216 Friesland Campina	16S ID	-		-	
163	<i>Staphylococcus intermedius</i>				RDCM 61.221	Unknown	-		-	

164	<i>Staphylococcus xylosus</i>				RDC 85	Unknown	-		-	
165	<i>Yersinia enterocolitica</i>				CIP 80.27T	Unknown	-		-	

¹Collection Institute Pasteur, Paris, France²Bio-Rad R&D Collection Marnes, Marnes, France³Culture Collection ADRIA Developpement, Quimper, France⁴Bio-Rad R&D Collection, Marnes, France⁵American Type Culture Collection, Manassas, VA⁶Bio-Rad Internal Collection, Steenvoorde, France⁷National Collection of Type Cultures, Salisbury, England⁸The Leibniz Institute DSMZ, Brunswick, Germany⁹Lavetan Reference Laboratory, Turnhout, Belgium¹⁰Belgian Coordinated Collections of Microorganisms, Ghent, Belgium¹¹Centre National de référence des Legionnelles, Lyon, France**Table 6: Summary of *Salmonella* Typhimurium Results (1)**

Level	With FDRS ¹	iQ-Check S. Typhimurium			Reference Method
		Presumptive	Without FDRS	Confirmed	
Raw chicken breast without skin					
Noninoculated	0/5	0/5	0/5	0/5	0/5
Low	8/20	8/20	8/20	8/20	8/20
High	5/5	5/5	5/5	5/5	5/5
Raw chicken breast with skin					
Noninoculated	0/5	0/5	0/5	0/5	0/5
Low	9/20	9/20	9/20	9/20	9/20
High	5/5	5/5	5/5	5/5	5/5
Raw chicken breast without skin containing 2% w/w salt					
Noninoculated	0/5	0/5	0/5	0/5	0/5
Low	8/20	8/20	8/20	8/20	8/20
High	5/5	5/5	5/5	5/5	5/5
Raw chicken thigh without skin					
Noninoculated	0/5	0/5	0/5	0/5	0/5
Low	10/20	10/20	10/20	10/20	10/20
High	5/5	5/5	5/5	5/5	5/5
Raw chicken thigh with skin					
Noninoculated	0/5	0/5	0/5	0/5	0/5
Low	7/20	7/20	7/20	7/20	7/20
High	5/5	5/5	5/5	5/5	5/5
Boot/Drag swabs					
Noninoculated	0/5	0/5	0/5	0/5	0/5
Low	8/20	8/20	8/20	8/20	7/20
High	5/5	5/5	5/5	5/5	5/5

¹FDRS = Free DNA Removal Solution

Table 7. iQ-Check S. Typhimurium Results – Candidate vs. Reference (1)

Matrix and Inoculum	MPN ^a / Test Portion	N ^b	Candidate			Reference			dPOD _C ^f	95% CI ^g
			x ^c	POD _C ^d	95% CI	x	POD _R ^e	95% CI		
Raw chicken breast without skin (25 g) <i>Salmonella</i> Typhimurium (ATCC BAA-215)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.55 (0.29, 0.93)	20	8	0.40	0.22, 0.61	8	0.40	0.22, 0.61	0.00	-0.13, 0.13
	2.58 (1.15, 5.78)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken breast with skin (25 g) <i>Salmonella</i> Typhimurium (ATCC 13311)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.61 (0.33, 1.04)	20	9	0.45	0.26, 0.66	9	0.45	0.26, 0.66	0.00	-0.13, 0.13
	1.97 (0.91, 4.27)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken breast without skin + 2% w/w salt (25 g) <i>Salmonella</i> Typhimurium (QL 11007-2)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.50 (0.25, 0.86)	20	8	0.40	0.22, 0.61	8	0.40	0.22, 0.61	0.00	-0.13, 0.13
	3.70 (1.52, 9.02)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken thigh without skin (25 g) <i>Salmonella</i> Typhimurium (ATCC 19585)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.70 (0.40, 1.14)	20	1 0	0.50	0.25, 0.75	10	0.50	0.25, 0.75	0.00	-0.13, 0.13
	1.97 (0.91, 4.27)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken thigh with skin (25 g) <i>Salmonella</i> Typhimurium (ATCC 14028)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.45 (0.22, 0.78)	20	7	0.35	0.18, 0.57	7	0.35	0.18, 0.57	0.00	-0.13, 0.13
	3.70 (1.52, 9.01)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Boot Swabs <i>Salmonella</i> Typhimurium (QL 011414.2)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	67	20	8	0.40	0.22, 0.61	7	0.35	0.18, 0.57	0.05	-0.23, 0.32
	180	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47

^aMPN = Most Probable Number is calculated using the LCF MPN calculator provided by AOAC RI, with 95% confidence interval; Test portions for boot swabs was determined by plating the inoculum

^bN = Number of test portions

^cx = Number of positive test portions

^dPOD_C = Candidate method confirmed positive outcomes divided by the total number of trials

^ePOD_R = Reference method confirmed positive outcomes divided by the total number of trials

^fdPOD_C = Difference between the confirmed candidate method and reference method confirmed POD values

^g95% CI = If the confidence interval of a dPOD does not contain zero, then the difference is statistically significant at the 5% level

Table 8. iQ-Check S. Typhimurium Results – Presumptive vs. Confirmed (1)

Matrix and Inoculum	MPN ^a /Test Portion	N ^b	Presumptive			Confirmed			dPOD _{CP} ^c	95% CI ^d
			x ^e	POD _{CP} ^d	95% CI	x	POD _R ^e	95% CI		
Raw chicken breast without skin (25 g) <i>Salmonella</i> Typhimurium (ATCC BAA-215)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.55 (0.29, 0.93)	20	8	0.40	0.22, 0.61	8	0.40	0.22, 0.61	0.00	-0.13, 0.13
	2.58 (1.15, 5.78)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken breast with skin (25 g) <i>Salmonella</i> Typhimurium (ATCC 13311)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.61 (0.33, 1.04)	20	9	0.45	0.26, 0.66	9	0.45	0.26, 0.66	0.00	-0.13, 0.13
	1.97 (0.91, 4.27)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken breast without skin + 2% w/w salt (25 g) <i>Salmonella</i> Typhimurium (QL 11007-2)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.50 (0.25, 0.86)	20	8	0.40	0.22, 0.61	8	0.40	0.22, 0.61	0.00	-0.13, 0.13
	3.70 (1.52, 9.02)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken thigh without skin (25 g) <i>Salmonella</i> Typhimurium (ATCC 19585)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.70 (0.40, 1.14)	20	10	0.50	0.25, 0.75	10	0.50	0.25, 0.75	0.00	-0.13, 0.13
	1.97 (0.91, 4.27)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Raw chicken thigh with skin (25 g) <i>Salmonella</i> Typhimurium (ATCC 14028)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	0.45 (0.22, 0.78)	20	7	0.35	0.18, 0.57	7	0.35	0.18, 0.57	0.00	-0.13, 0.13
	3.70 (1.52, 9.01)	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47
Boot Swabs <i>Salmonella</i> Typhimurium (QL 011414.2)	-	5	0	0.00	0.00, 0.43	0	0.00	0.00, 0.43	0.00	-0.47, 0.47
	67	20	8	0.40	0.22, 0.61	8	0.40	0.22, 0.61	0.00	-0.13, 0.13
	180	5	5	1.00	0.57, 1.00	5	1.00	0.57, 1.00	0.00	-0.47, 0.47

^aMPN = Most Probable Number is calculated using the LCF MPN calculator provided by AOAC RI, with 95% confidence interval; Test portions for boot swabs was determined by plating the inoculum

^bN = Number of test portions

^cx = Number of positive test portions

^dPOD_{CP} = Candidate method presumptive positive outcomes divided by the total number of trials

^ePOD_R = Candidate method confirmed positive outcomes divided by the total number of trials

^fdPOD_{CP} = Difference between the candidate method presumptive and confirmed POD values

^g95% CI = If the confidence interval of a dPOD does not contain zero, then the difference is statistically significant at the 5% level

^hResults between traditional and alternative confirmation were identical

DISCUSSION OF THE MODIFICATION STUDY APPROVED JANUARY 2023 (5)

The new CFX Opus Deepwell instrument delivers the same performance as the current CFX96 Touch Deep Well instrument but with a more modern design and cloud capabilities. The improved stability of the thermal block ensures a more uniform thermal protocol. The CFX Manager Software, IDE v 3.1 brings the same performance, algorithm, and interpretation as the current CFX Manager Software, IDE v 3.0 with the only change being compatibility to both CFX96 Touch Deep Well and CFX Opus Deepwell instruments.

REFERENCES CITED

1. Clark, M., Pourniaire, JP., Pierre, S., and Quiring, C., Validation for the iQ-Check S. Enteritidis and S. Typhimurium Real-Time PCR Methods, iQ-Check S. Enteritidis, AOAC Performance Tested MethodsSM certification number 081903.
2. U.S. Department of Agriculture Food Safety and Inspection Service (2018) Microbiology Laboratory Guidebook, Revision 4.09 *Isolation and Identification of Salmonella from Meat, Poultry, Pasteurized Egg, and Siluriformes (Fish) Products and Carcass and Environmental Sponges* (Accessed December 2018)
3. FDA, *Environmental Sampling and Detection of Salmonella in Poultry Houses*.
<https://www.fda.gov/food/foodscienceresearch/laboratorymethods/ucm114716.htm> (Accessed December 2018)
4. iQ-Check Salmonella Enteritidis and iQ-Check Salmonella Typhimurium User Guides, December 2018
5. Clark, M., Validation of the Group Modification for the Addition of the CFX Opus Deepwell Real-Time PCR Instrument and CFX Manager Software, IDE v3.1, AOAC Performance Tested MethodsSM certification number 081904. Modification approved January 2023.